

$T > 0$ Ensemble State Density Functional Theory Revisited

Helmut Eschrig*

IFW Dresden, PO Box 270116, D-0111171 Dresden, Germany

A logical foundation of equilibrium state density functional theory in a Kohn-Sham type formulation is presented on the basis of Mermin's treatment of the grand canonical state. It is simpler and more satisfactory compared to the usual derivation of ground state theory, and free of remaining open points of the latter. It may in particular be relevant with respect to cases of spontaneous symmetry breaking like non-collinear magnetism and orbital order.

PACS numbers: 05.30.Fk, 31.15.ec, 71.15.Mb

I. INTRODUCTION

Modern ground state density functional theory (DFT) for an inhomogeneous system of identical particles, having early roots in the work of Thomas and Fermi, was pioneered by the seminal papers by Hohenberg, Kohn and Sham.^{1,2} It was later generalized by the constrained search concept of Levy³ and finally put on a mathematically rigorous basis of functional Legendre transforms by Lieb.⁴ Meanwhile, DFT for the quasi-particle self-energy labeled by acronyms like GW or LDA+DMFT, and time-dependent DFT for dynamical processes, related to Keldysh Green's functions, were developed, and all these theories laid the ground for enormously successful model approaches (by use of model functionals) to simulation of molecules and solids of any complexity.

A generalization of the ground state DFT to thermodynamic equilibrium states by Mermin⁵ which appeared shortly after the work of Hohenberg, Kohn and Sham, has been mentioned from time to time,^{6,7} but otherwise has been largely ignored so far.

As it sometimes happens,⁸ a formalism for temperature $T = 0$ need not be equivalent to that for $T \downarrow 0$ which latter case is always the relevant case in physics. Although ground state DFT is largely settled now, some uneasy feeling remains in connection with the density functional failing to be differentiable in some cases (where $n_{ss'} \mapsto v_{ss'}$ is not unique), notably in spin DFT.^{9,10} In Ref. 9 it is argued that these problems may reduce to the ordinary well understood gap problem, now for the spin subsystems separately, if one restricts consideration to homogeneous external magnetic fields only. This might seem a reasonable restriction since applied fields in laboratory can hardly vary over microscopic distances. However, in the very topical cases of spontaneous symmetry breaking with respect to the interplay of non-collinear magnetism with orbital order, in a statistical treatment one has to resort to the trick of Bogolubov's quasi-means by applying a suitable infinitesimal symmetry-breaking external field, otherwise statistical ensembles would not reproduce the broken symmetry. In the just mentioned cases this implies a microscopically inhomogeneous field, and one would like to rely on a situation where everything is fine at least in an infinitesimal vicinity of such a symmetry-breaking field. The good news is that this

is indeed the case and the needed functional derivatives always exist for $T > 0$. This is shown with the help of Mermin's approach in the sequel. DFT is a rigorous theory for volume $V < \infty$ and for temperature $T > 0$. For $V = +\infty$ a ground state wave function may not exist and for $T = 0$ the functional derivative of the density functional may not exist. The theory then may be applied for $V \rightarrow \infty$ (as is routinely done with refinement of the grid in $\mathbf{k}$ -space; a discrete regular $\mathbf{k}$ grid means periodic boundary conditions with a finite periodicity volume) and for $T \downarrow 0$. (Contrary to the case of adiabatic molecular dynamics⁷ the temperature of a finite system in an equilibrium state has a well defined meaning in the average over states the system may be in after it had been for a long time in contact with a large thermal bath.)

II. SUBTLETIES OF HOHENBERG-KOHN-SHAM THEORY

Originally¹ DFT was built for systems in an (arbitrarily large) box of finite volume which conveniently can be replaced by periodic boundary conditions meaning to treat the position space of the particles as a three-torus $\mathbb{T}^3$ of finite volume $|\mathbb{T}^3|$, for instance $x \equiv x + L$, $y \equiv y + L$, $z \equiv z + L$, $|\mathbb{T}^3| = L^3$.

Let the Hamiltonian be

$$\hat{H} = \hat{T} + \hat{W} + \hat{V}, \quad (1)$$

$$\hat{T} = \frac{\hbar^2}{2m} \int \nabla \hat{\psi}^\dagger(x) \nabla \hat{\psi}(x) dx, \quad (2)$$

$$\hat{W} = \int \hat{\psi}^\dagger(x) \hat{\psi}^\dagger(x') w(\mathbf{r}, \mathbf{r}') \hat{\psi}(x') \hat{\psi}(x) dx' dx, \quad (3)$$

$$\hat{V} = \int \hat{\psi}^\dagger(\mathbf{r}, s) v(\mathbf{r}) \hat{\psi}(\mathbf{r}, s) dx, \quad (4)$$

where $\hat{\psi}(x)$, $x = (\mathbf{r}, s)$ with particle position $\mathbf{r}$ and spin variable s , is the field operator of the particle field and $\int dx = \sum_s \int d^3r$.

Then, given any particle number N , a normalized ground state (GS) wavefunction (WF) $\Psi(x_1, \dots, x_N)$ and

a GS density $n(\mathbf{r})$ exist for any reasonable external potential $v(\mathbf{r})$ and for any non-negative pair interaction $w(\mathbf{r}_i, \mathbf{r}_j)$. As is now standard,⁴ one allows for all potentials with the only condition that $\int_{\mathbb{T}^3} |v|^{3/2} d^3r < \infty$, that is, $v \in \mathbf{L}^{3/2}(\mathbb{T}^3)$. Potentials of arrays of finitely many point charges in the $\mathbb{T}^3$ belong to this space^{4,11}, and the Hamiltonian $\hat{H}^0 = \hat{T} + \hat{V}$ of interaction-free fermions is bounded below for any such potential. Then, this also holds true for Hamiltonians (1), if $w(\mathbf{r}, \mathbf{r}') \geq 0$. Since the space $\mathbb{T}^3$ has finite volume, all considered Hamiltonians have discrete spectra with at most finite degrees of level degeneracy. (Lieb allowed the position space to be the real vector space $\mathbb{R}^3$ of infinite volume which caused many problems with the continuous part of the spectrum of Hamiltonians. He then had to restrict $n \in \mathbf{L}^3(\mathbb{R}^3) \cap \mathbf{L}^1(\mathbb{R}^3)$ since the density must integrate to a finite particle number N over the infinite space $\mathbb{R}^3$. This led him allow for potentials $v \in \mathbf{L}^{3/2}(\mathbb{R}^3) + \mathbf{L}^\infty(\mathbb{R}^3)$. In the three-torus every function $n \in \mathbf{L}^3(\mathbb{T}^3)$ may be normalized to integrate to a given N .)

The lemma by Hohenberg and Kohn¹ states the unique mapping $n \mapsto v$ from ground state densities (degenerate ground states allowed⁴) to external potentials, on which basis the Hohenberg-Kohn density functional

$$F_{\text{HK}}[n] = E[v[n], N] - \langle n, v[n] \rangle \quad (5)$$

is defined for any ground state density $n \in \mathcal{A}_N$:

$$\mathcal{A}_N = \{n \text{ coming from an } N\text{-particle GS-WF}\}. \quad (6)$$

In (5) $v[n]$ means the potential causing a WF-GS density n , and henceforth we use the notation of linear functionals

$$\langle n, v \rangle = \int_{\mathbb{T}^3} n(\mathbf{r}) v(\mathbf{r}) d^3r. \quad (7)$$

As is easily seen, $\langle n, v \rangle$ cancels an equal term in the GS energy $E[v[n], N]$, so that F_{HK} does not any more depend on $v[n]$. The functional $F = F_{\text{HK}}$ might be used in the variational principle by Hohenberg and Kohn

$$E[v, N] = \min_n \{F[n] + \langle v, n \rangle \mid \langle 1, n \rangle = N\} \quad (8)$$

where $\{A|B\}$ means a set of elements A with property B . The crucial point for the possibility to solve this problem with the help of Euler's equation is the knowledge of the variational domain for n and the existence of the functional derivative of F . Would the derivative of F at the minimizing density $n_{\min}$ exist, it would be $\delta F[n_{\min}]/\delta n = -v + \mu$, where μ is the Lagrange multiplier for the constraint in (8). For F_{HK} , unfortunately neither the domain of definition $\mathcal{A}_N$ is explicitly known nor is anything known about the existence of the functional derivative. We know that $\mathcal{A}_N \subset \mathbf{L}^3(\mathbb{T}^3)$, but Lieb has shown⁴ that $\mathcal{A}_N$ is not convex: There are densities $n = \sum_i c_i n_i$, $c_i \geq 0$, $\sum_i c_i = 1$, which are not in $\mathcal{A}_N$ while the n_i all are in $\mathcal{A}_N$. This is, why nowadays more general definitions of $F[n]$ are used.

As the theory can be build for any reasonable pair interaction w , the interaction-free case $w = 0$ is of some help. Further on this case will be denoted by a superscript 0, but the corresponding density functionals F^0 will as usually be denoted by T since they obviously reduce to the kinetic energy of an interaction-free system with GS density n . In this case, an alternative to (5) is the density matrix (DM) functional

$$T_{\text{DM}}[n] = \min_{\substack{\{p_k, \varphi_k\} \\ 0 \leq p_k \leq 1 \\ (\varphi_k | \varphi_{k'}) = \delta_{kk'}}} \left\{ \sum_k p_k (\varphi_k | \hat{t} | \varphi_k) \mid \sum_k p_k |\varphi_k|^2 = n \right\} \quad (9)$$

with $\hat{t} = -\hbar^2 \Delta / 2m$. For any N -particle DM state (not only GS),

$$\begin{aligned} \mathcal{T}[\{p_k, \varphi_k\}] &= \sum_k p_k (\varphi_k | \hat{t} | \varphi_k), \\ 0 \leq p_k \leq 1, \quad \sum_k p_k &= N, \quad (\varphi_k | \varphi_{k'}) = \delta_{kk'}, \end{aligned} \quad (10)$$

is the general expression of the kinetic energy (with the set of orthonormal orbitals φ_k depending on the state). Given a potential v , the orbitals with $(\hat{t} + v)\varphi_k = \varphi_k \varepsilon_k$ and occupation numbers $p_k = 1$ for $\varepsilon_k < \varepsilon_N$, $p_k = 0$ for $\varepsilon_k > \varepsilon_N$ minimize (9) for the corresponding GS density $n \in \mathcal{A}_{\text{DM}, N}^0$. The GS is unique if the highest occupied level ε_N is not degenerate. The GSs and their energies are obtained from the Kohn-Sham (KS) variational principle

$$E^0[v, N] = \min_{\substack{\{p_k, \varphi_k\} \\ 0 \leq p_k \leq 1, \sum p_k = N \\ (\varphi_k | \varphi_{k'}) = \delta_{kk'}}} \left\{ \mathcal{T}[\{p_k, \varphi_k\}] + \sum_{k=1}^N (\varphi_k | v | \varphi_k) \right\} \quad (11)$$

while (9) is defined for the density n of any N -particle state, that is,⁴ on the domain

$$\mathcal{J}_N = \{n \mid n(\mathbf{r}) \geq 0, \nabla(n^{1/2}) \in \mathbf{L}^2(\mathbb{T}^3), \langle 1, n \rangle = N\} \quad (12)$$

which is a convex subset of $\mathbf{L}^3(\mathbb{T}^3)$. (It has been shown⁴ that a minimum (9) exists for every $n \in \mathcal{J}_N$.)

Let for the sake of simplicity v have a non-degenerate GS (which is a single determinant of orbitals in this case) with density n . It is easily seen that the most general variation permitted by (9) is a linear combination of $\{\delta\varphi_k = \lambda_k \varphi_{l_k}\}$, $\varepsilon_k \leq \varepsilon_N$, $\varepsilon_{l_k} > \varepsilon_N$, $\lambda_k \rightarrow 0$. To lowest order in the λ_k it yields $\delta T_{\text{DM}} = 2\text{Re} \sum_k \lambda_k (\varphi_{l_k} | \hat{t} | \varphi_k) = 2\text{Re} \sum_k \lambda_k (\varphi_{l_k} | v | \varphi_k) = \langle v, \delta n \rangle$. These variations reach every single determinant state in a neighborhood of the considered GS (with respect to the $\mathbf{H}^1(\mathbb{T}^{3N})$ -norm $\|\Psi\|^2 = \int (|\Psi|^2 + |\nabla \Psi|^2) d^3N r$). Since⁴ single determinant states map onto $\mathcal{J}_N$, the corresponding δn is a general variation in a neighborhood of n in $\mathcal{J}_N$ (relative to the $\mathbf{L}^3(\mathbb{T}^3)$ -norm), and hence the functional derivative of the convex functional T_{DM} exists at n as a (Frechet) derivative in $\mathcal{J}_N$ and equals v . The argument can be

generalized to the case of a degenerate GS, that is, for all $n \in \mathcal{A}_{\text{DM},N}^0$.

Now, let $n \in \mathcal{J}_N \setminus \mathcal{A}_{\text{DM},N}^0$. Such densities exist, for instance densities having nodes cannot be in $\mathcal{A}_{\text{DM},N}^0$. Assume that the derivative of T_{DM} exists at that n . This means that there is some u with $\delta T_{\text{DM}} = \langle u, \delta n \rangle$ for all permitted δn . Since T_{DM} was shown⁴ to be convex, the assumption implies that n minimizes $T_{\text{DM}}[n'] - \langle u, n' \rangle$ and hence is a GS density to the potential $-u$ in contradiction to the presupposition. T_{DM} has nowhere outside of $\mathcal{A}_{\text{DM},N}^0$ a functional derivative.

KS theory in the interacting case $w > 0$ now uses the splitting

$$F[n] = T_{\text{DM}}[n] + E_{\text{H}}[n] + E_{\text{XC}}[n], \quad (13)$$

which defines the density functional $E_{\text{XC}}[n]$ through the preceding ones. While this definition is correct, nothing can be said on the existence of the functional derivative of $E_{\text{XC}}[n]$ for GS densities $n \in \mathcal{A}_{\text{DM},N}$ since we do not know the sets $\mathcal{A}_{\text{DM},N}^0$ and $\mathcal{A}_{\text{DM},N}$ and cannot assume $\mathcal{A}_{\text{DM},N} \subset \mathcal{A}_{\text{DM},N}^0$. Even though F has a functional derivative for $n \in \mathcal{A}_{\text{DM},N}$ (see below), $E_{\text{XC}}[n]$ can only have one there if T_{DM} has one.

Like in (11), with $n = \sum_k p_k |\varphi_k|^2$ a KS variational principle is set out with the KS equation as the corresponding Euler equation. Now one may assume

$$\begin{aligned} T_{\text{DM}}[n] + E_{\text{XC}}[n] = \\ = \min_{\substack{\{p_k, \varphi_k\} \\ 0 \leq p_k \leq 1 \\ (\varphi_k | \varphi_{k'}) = \delta_{kk'} \\ \left| \sum_{k=1}^N p_k |\varphi_k|^2 = n \right|}} \left\{ \sum_{k=1}^N p_k (\varphi_k | \hat{t} | \varphi_k) + \mathcal{E}_{\text{XC}}[\{p_k, \varphi_k\}] \right\} \end{aligned} \quad (14)$$

leading to a KS equation with a (non-linear) exchange and correlation potential operator

$$\frac{1}{p_k} \frac{\delta \mathcal{E}_{\text{XC}}}{\delta \varphi_k^*(\mathbf{r})} = \hat{v}_{\text{XC}} \varphi_k(\mathbf{r}), \quad (\hat{v}_{\text{XC}} \varphi_k)^* = \hat{v}_{\text{XC}} \varphi_k^*, \quad (15)$$

if one assumes that $\mathcal{E}_{\text{XC}}$ depends on p_k and on a Hermitian form of the φ_k only. Like in (9), the right hand side of (14) might exist (not proven so far; only $\mathcal{E}_{\text{X}}[\{p_k, \varphi_k\}]$ is a simple explicitly known expression). By inserting (14) together with $n = \sum_k p_k |\varphi_k|^2$ into (13) it is seen that $\mathcal{E}_{\text{XC}}$ (if it exists) has derivatives with respect to p_k and φ_k since all the other terms in the equation have them (even where the left hand side of (14) has no derivative with respect to n). Hence, $\hat{v}_{\text{XC}}$ will in general not be a local potential function, it may in particular be orbital dependent ($\hat{v}_{\text{X}}$ is non-local, its orbital dependence is canceled by inclusion of the orbital dependent self-interaction in both v_{H} and v_{X} ; the model XC potentials with partial self-interaction correction or in LDA+ U models are non-local and orbital dependent). Would v_{XC} exist as a local potential, then the KS equation would always yield a solution $n \in \mathcal{A}_{\text{DM},N}^0$ and hence there would

be $\mathcal{A}_{\text{DM},N} \subset \mathcal{A}_{\text{DM},N}^0$ which can by no means be taken for granted.

The only density functional F for which the issue of the existence of the functional derivative can be addressed in general is the Legendre transform⁴

$$F[n] = \sup_{v \in X^*} \{E[v, N] - \langle n, v \rangle\} \quad (16)$$

for both cases, $w = 0$ and $w \neq 0$. It is convex and defined on the whole functional space X (it may take on the value $+\infty$ in part of X), and if, given n , there exists a unique maximizing v , then this is the functional derivative of $F[n]$. Since v is indeed up to a constant uniquely determined by any GS n , the functional derivative of this F exists at least for $n \in \mathcal{A}_{\text{DM},N}$ as a derivative ('gradient', more precisely Frechet derivative) in the hyperplane $\{n \in X | \langle 1, n \rangle = N\}$.

Less clear is the situation in spin DFT.⁹ Now, also F need not have a derivative for GS densities.

III. A FEW ESSENTIALS ON LEGENDRE TRANSFORMS

Let $X = X^{**}$ and X^* be two mutually dual functional spaces, that is, X^* comprises all norm-continuous linear functions on X and vice versa. $((\mathbb{R}^N)^*)^* = \mathbb{R}^N$ is the space of all gradient vectors to functions on $\mathbb{R}^N$; $(\mathbf{L}^3(\mathbb{T}^3))^* = \mathbf{L}^{3/2}(\mathbb{T}^3)$ and vice versa.) The Legendre transform $f^*(n)$, $n \in X$ of a function $f(v)$, $v \in X^*$ is defined as

$$f^*(n) = \sup_{v \in X^*} \{\langle n, v \rangle - f(v)\}. \quad (17)$$

A second Legendre transformation yields

$$f^{**}(v) = \sup_{n \in X} \{\langle v, n \rangle - f^*(n)\}. \quad (18)$$

All we need is

1. $f^*(n)$ is a convex function of n , no matter what $f(v)$ is; if $f(v)$ is convex, then $f^{**}(v) = f(v)$; in general $f^{**}(v) \leq f(v)$.
2. $f(v) + f^*(n) \geq \langle v, n \rangle$; if, for convex f , $f(v) + f^*(n) = \langle v, n \rangle$, then $v \in \partial f^*(n)$ and $n \in \partial f(v)$.

In the second statement $\partial f(v)$ is the subdifferential of the convex function f at point v : the set of all linear functions $\langle n, v' \rangle$ so that $f(v') \geq f(v) + \langle n, (v' - v) \rangle$ for all $v' \in X^*$. If this set consists of a single linear function only, then this linear function is the (total) differential $df(v)$, that is, n is the derivative of f at v .

To elucidate these properties one may consider convex functions of one real variable, $f(N)$ and $f^*(\mu)$ (see e.g. Fig. 11 of Ref. 11). Put a supporting tangent to the graph of f at point N (a line having the common point $(N, f(N))$ with the graph of the function and being nowhere above). The tangent has a slope μ . The

sign carrying distance from the intersection point of this line with the f -axis to the coordinate origin is $f^*(\mu)$. If f has a derivative at N , then its value is μ . It is easily seen that, if the derivative of f jumps at N , then there is a (closed) interval $[\mu_1, \mu_2]$ from the left derivative μ_1 to the right derivative μ_2 (μ_1 may be $-\infty$ or μ_2 may be $+\infty$), and $f^*(\mu)$ is *linear* on this interval, the interval being the subdifferential $\partial f(N)$. Inversely, if the convex function f is not strictly convex, but has a linear dependence on some interval with slope μ , then the derivative of f^* jumps at that μ .

This simple geometric picture readily transfers to the general case: take a tangent hyperplane $f(n_0) + \langle v, (n - n_0) \rangle$ supporting the graph of $f(n)$ at some n_0 . The distance from its intersection point with the f -axis to the origin is $f^*(v)$. If the derivative of f jumps at some n (and hence does not exist there), then there is a convex domain in v -space on which $f^*(v)$ is linearly depending on v , and vice versa.

If the GS wave function is independent of some potential change δv called a ‘phantom’ potential perturbation in Ref. 10, then the GS density n does also not change and the GS energy has a linear dependence $\text{const.} + \langle \delta v, n \rangle$. Consequently, the functional derivative of $F[n]$ defined by (16) does not exist at that n . This is precisely the role of ‘phantom’ potential perturbations in DFT.

IV. UNIQUE MAPPINGS FOR $T > 0$

We now move to temperature $T > 0$ and to grand canonical states. We also generalize to spin DFT and allow for external magnetic fields coupling to the particle spin but not to its charge (diamagnetic couplings as usually in spin DFT are neglected). Consider a system of identical particles in an external field $v_{ss'}(\mathbf{r})$. Let the system be confined in a large box, or, placed in a large three-torus equivalent to periodic boundary conditions (regular $\mathbf{k}$ -grid). Let the Hamiltonian be that of (1-3), but (4) generalized to

$$\hat{V} = \sum_{ss'} \int \hat{\psi}^\dagger(\mathbf{r}, s) v_{ss'}(\mathbf{r}) \hat{\psi}(\mathbf{r}, s') d^3r, \quad (19)$$

The particle number operator is $\hat{N} = \int \hat{\psi}^\dagger(x) \hat{\psi}(x) dx$ so that $\hat{H} - \mu \hat{N}$ depends on the combination $v - \mu = v_{ss'}(\mathbf{r}) - \mu \delta_{ss'}$ only.

Fix the temperature $\beta = 1/kT$, the chemical potential μ and the external potential v . Then, the grand canonical state is

$$\rho_\beta[v - \mu] = \frac{e^{-\beta(\hat{H} - \mu \hat{N})}}{\text{tr } e^{-\beta(\hat{H} - \mu \hat{N})}}. \quad (20)$$

If $\rho > 0$, $\text{tr } \rho = 1$, is *any* state (density matrix), then $\text{tr } \rho(\hat{V} - \mu \hat{N}) = \int (v - \mu) n[\rho] dx$ with the particle (spin) density

$$n[\rho] = n_{ss'}(\mathbf{r}) = \text{tr } \rho \hat{\psi}(\mathbf{r}, s) \hat{\psi}^\dagger(\mathbf{r}, s'). \quad (21)$$

In the following tr will always mean the trace in the Fock space of the $\hat{\psi}$. Also, the natural abbreviation

$$\sum_{ss'} \int (v_{ss'}(\mathbf{r}) - \mu \delta_{ss'}) n_{s's}(\mathbf{r}) d^3r = \langle (v - \mu), n \rangle \quad (22)$$

will be used.

Now, fix the particle interaction w and, following Mermin⁵ (we try carefully to trace functional dependencies and in doing so slightly deviate from Mermin’s notation), consider for various external potentials v the functionals

$$\Omega_v[\rho] = \text{tr } \rho \left(\hat{H} - \mu \hat{N} + \frac{1}{\beta} \ln \rho \right). \quad (23)$$

As easily seen by direct substitution of (20), the grand canonical potential $\Omega_\beta[v - \mu]$ is obtained as

$$\Omega_\beta[v - \mu] = -\frac{1}{\beta} \ln \text{tr } e^{-\beta(\hat{H} - \mu \hat{N})} = \Omega_v[\rho_\beta[v - \mu]]. \quad (24)$$

Moreover, as shown in Ref. 5, for any $\rho > 0$, $\text{tr } \rho = 1$, it holds that

$$\Omega_v[\rho] > \Omega_v[\rho_\beta[v - \mu]] = \Omega_\beta[v - \mu] \text{ for } \rho \neq \rho_\beta[v - \mu]. \quad (25)$$

In Mermin’s approach, this inequality replaces the corresponding ground state energy property. It immediately implies that $\Omega_\beta[v - \mu] = \min_\rho \Omega_v[\rho]$ is concave in v by the simple reasoning (we write in short v_i for $v_i - \mu_i$)

$$\begin{aligned} \Omega_\beta[\alpha v_1 + (1 - \alpha)v_2] &= \\ &= \min_\rho \text{tr } \rho \left(\alpha \hat{H}_1 + (1 - \alpha) \hat{H}_2 + \frac{1}{\beta} \ln \rho \right) \geq \\ &\geq \alpha \min_{\rho_1} \text{tr } \rho_1 \left(\hat{H}_1 + \frac{1}{\beta} \ln \rho_1 \right) + \\ &\quad (1 - \alpha) \min_{\rho_2} \text{tr } \rho_2 \left(\hat{H}_2 + \frac{1}{\beta} \ln \rho_2 \right) = \\ &= \alpha \Omega_\beta[v_1] + (1 - \alpha) \Omega_\beta[v_2], \quad 0 \leq \alpha \leq 1, \end{aligned} \quad (26)$$

because a joint minimum of a sum cannot be below the sum of the independent minima of the items.

As another advantage over the standard zero temperature theory, it follows immediately from (20) and (21) that the mappings $(v - \mu) \mapsto \rho_\beta \mapsto n$ are unique. There is no problem with degenerate states since degenerate states automatically get equal statistical weight in ρ_β of (20). However, as usual spontaneous symmetry breaking is not covered by this statistical approach; it has to be treated by an infinitesimal symmetry breaking external potential v in the spirit of Bogolubov’s quasi-means in Statistical Physics. Nevertheless, by virtue of (25) which also holds in the spin case, in the standard way Mermin proved the analogue of the Hohenberg-Kohn lemma: $n \mapsto (v - \mu)$ is unique for any n coming from a grand canonical ensemble at temperature $1/k\beta$. In summary,

there are the unique mappings

$$\begin{array}{ccc} (v - \mu) & \longleftarrow & n \\ \downarrow & & \uparrow \\ \text{grand canonical } \rho_\beta & & \end{array} \quad (27)$$

On the functional domain (which may depend on $\hat{W}$)

$$D_\beta = \{n \text{ coming from some } \rho_\beta, \beta \text{ fixed}\} \quad (28)$$

one can write $(v - \mu)_\beta[n]$ and also $\hat{H}_\beta[n]$ and $\rho_\beta[n]$, as well as $n_\beta[v - \mu] = n[\rho_\beta[v - \mu]]$ on the domain of admissible potentials v . (Denoting the distinct functions $\rho_\beta[n]$ and $\rho_\beta[v]$ by the same symbol ρ_β will cause no confusion.)

Moreover, from the unique dependence of $v - \mu$ on ρ_β it follows now also that $\Omega_v[\rho]$ for different $v_i - \mu_i$ is minimized by different ρ_i , and hence the inequality in (26) is sharpened into a strict inequality: $\Omega_\beta[v]$ is strictly concave. If equality would hold in (26), this would imply that the minimizing ρ is also a minimizing ρ_1 for v_1 and a minimizing ρ_2 for v_2 . This is the principal difference from the $T = 0$ theory where $E[v, N]$ is not always strictly concave in $v_{ss'}$ and is never strictly convex in N .

V. THE DENSITY FUNCTIONAL

As was already said, for electron systems it is well justified to allow for all potentials

$$v - \mu \in \mathbf{L}^{3/2}(\mathbb{T}^3) = X^* \quad (29)$$

for which the integral $\int_{\mathbb{T}^3} |v - \mu|^{3/2} d^3r$ over the three-torus (of finite volume) is finite. Recall that the Hamiltonian $\hat{H}^0 = \hat{T} + \hat{V}$ of interaction-free fermions is bounded below for any such potential, that this also holds true for Hamiltonians (1-3, 19), if $w(\mathbf{r}, \mathbf{r}') \geq 0$, and that, since the space $\mathbb{T}^3$ has finite volume, all considered Hamiltonians have discrete spectra with at most finite degrees of level degeneracy. Then, $\Omega_\beta[v - \mu]$ of (24) is well defined on X^* and smooth in the norm topology.

In view of the concavity of $\Omega_\beta[v]$, introduce the Legendre transform^{4,11} of $-\Omega_\beta[v]$ as $\tilde{F}_\beta[-n]$:

$$F_\beta[n] = \tilde{F}_\beta[-n] = \sup_v \left\{ -\langle n, v \rangle + \Omega_\beta[v] \right\} \quad (30)$$

which as a Legendre transform is a convex functional of $-n$ (or likewise of n), the dual variable to v : $n \in X^{**} = X = \mathbf{L}^3(\mathbb{T}^3)$.

Since the functional space X is reflexive, $\mathbf{L}^3(\mathbb{T}^3) = (\mathbf{L}^3(\mathbb{T}^3))^{**}$, the Legendre back transformation from (30), $-\Omega_\beta[v] = \sup_n \{ -\langle v, n \rangle - \tilde{F}_\beta[-n] \}$ or,

$$\Omega_\beta[v] = \inf_n \left\{ F_\beta[n] + \langle v, n \rangle \right\} \quad (31)$$

represents the generalized Hohenberg-Kohn theorem (where equality holds since $\Omega_\beta[v]$ is concave in v). The

chemical potential μ is further on put to zero which simply means that single particle energies and potentials are measured from the chemical potential.

For any density $n \in D_\beta$ from (28), in analogy to the original Hohenberg-Kohn functional one may define

$$\Phi_\beta[n] = \Omega_v[\rho_\beta[n]] - \langle v, n \rangle, \quad n \in D_\beta \quad (32)$$

and, from (25), have

$$\Omega_\beta[v] = \min_{n \in D_\beta} \left\{ \Phi_\beta[n] + \langle v, n \rangle \right\} \quad (33)$$

since in view of (27) any $n \neq n[v]$ refers to $\rho_\beta[n] \neq \rho_\beta[n[v]]$. From (31) and (33) one infers that $F_\beta[n] = \Phi_\beta[n]$ for all $n \in D_\beta$, and that the infimum of (31) is always a minimum with minimizing density $n_\beta[v]$: $\Phi_\beta[n_\beta[v]] + \langle v, n_\beta[v] \rangle = \Omega_\beta[v] \leq F_\beta[n_\beta[v]] + \langle v, n_\beta[v] \rangle$, hence $\Phi_\beta[n] \leq F_\beta[n]$ for $n \in D_\beta$, and by interchanging the role of (31) and (33) in the argument the opposite inequality is obtained.

Moreover, for $-\Omega_\beta[v]$ and $\tilde{F}_\beta[-n]$ like in general for any pair of mutual Legendre transforms it holds that $\tilde{F}_\beta[-n] - \Omega_\beta[v] = -\langle n, v \rangle$ implies $v \in \partial \tilde{F}_\beta[-n]$ and $-n \in \partial(-\Omega_\beta[v])$, where $\partial \tilde{F}_\beta[-n]$ means the subdifferential on $\tilde{F}_\beta$ at point $-n$ and $\partial(-\Omega_\beta[v])$ means the subdifferential on $-\Omega_\beta$ at point v . Since $d\Omega_\beta[v] = \ln \text{tr}(e^{-\beta \hat{H}} d\hat{V})$ as easily seen from the definition of $\Omega_\beta[v]$, its first (Frechet) derivative (for finite volume $|\mathbb{T}^3|$) always exists so that its subdifferential contains only this one 'gradient'. Now, the reasoning after (33) yields $F_\beta[n_\beta[v]] - \Omega_\beta[v] = -\langle n_\beta[v], v \rangle$ and hence

$$\frac{\delta \Omega_\beta}{\delta v} = n_\beta[v]. \quad (34)$$

As $n_\beta[v]$ is a one-one mapping $X^* \leftrightarrow D_\beta$, for $n \in D_\beta$ one has inversely $v_\beta[n]$ and $F_\beta[n] - \Omega_\beta[v_\beta[n]] = -\langle n, v_\beta[n] \rangle$ implying

$$\frac{\delta F_\beta}{\delta n} = -v_\beta[n], \quad n \in D_\beta. \quad (35)$$

Note that while (35) holds for $n \in D_\beta \subset X$, the derivative $\delta/\delta n$ on the left hand side is taken in X , that is, for any $\delta n \in X$ with $\|\delta n\|$ small enough.

From the strict concavity and continuous differentiability of $\Omega_\beta[v]$ the differentiability of $F_\beta[n]$ at every point $n \in D_\beta$ follows, that is, at every density n thermodynamically corresponding to some v at temperature $(k\beta)^{-1}$. Like in the $T = 0$ theory^{4,11}, (30) yields that $F_\beta[n]$ jumps to $+\infty$ if $n < 0$ for an x -domain of non-zero measure: Take $v = c > 0$ for some domain where $n < 0$ and $v = 0 (= \mu)$ everywhere else. This v is admissible for arbitrary large c and $\hat{H}[v]$ is bounded below for such a v . Hence, $\Omega_\beta[v]$ is also bounded below and, as easily seen, the supremum (30) is obtained for $c \rightarrow \infty$ to be $+\infty$. Assume now that $n_\beta(x_0) = 0$ for some x_0 . Since any $n_\beta \in D_\beta$ is continuous in x (any solution of the many-particle Schrödinger equation is continuous), there is always $\delta n \in X$, $\delta n(x_0) > 0$, so that $n_\beta(x) - \epsilon \delta n(x)$ would

be negative in a neighborhood of x_0 of non-zero measure for arbitrarily small $|\epsilon|$ and the functional derivative (35) would not exist for that n_β . Thus, the result (35) also implies $n_\beta(x) > 0$ everywhere for $T > 0$. (See also next section.)

VI. INTERACTION-FREE PARTICLES AND BEYOND

As is well known from Statistical Physics¹², in a non-interacting particle system the particles in a single particle quantum state $|\varphi_k\rangle$ may be treated as an independent subsystem even of a quantum ensemble with exchange symmetry. The corresponding statistical fermionic state is

$$\rho_k = | \rangle (1 - p_k) \langle | + | \varphi_k \rangle p_k \langle \varphi_k | \quad (36)$$

with occupation p_k of the orbital φ_k and $| \rangle$ as the vacuum state. Accordingly we define⁷

$$\begin{aligned} \mathcal{T}_\beta[p_k, \varphi_k] &= \text{tr } \rho_k \left(\hat{T} + \frac{1}{\beta} \ln \rho_k \right) = \\ &= -p_k (\varphi_k^* | \hat{t} | \varphi_k) + \\ &+ \frac{1}{\beta} (p_k \ln p_k + (1 - p_k) \ln(1 - p_k)) \end{aligned} \quad (37)$$

and the density functional

$$T_\beta[n] = \min_{\substack{\{p_k, \varphi_k\} \\ 0 \leq p_k \leq 1 \\ (\varphi_k | \varphi_{k'}) = \delta_{kk'}}} \left\{ \sum_k \mathcal{T}_\beta[p_k, \varphi_k] \mid \sum_k p_k |\varphi_k|^2 = n \right\} \quad (38)$$

where the minimum taken over all orthonormal orbitals and orbital occupations which yield a given n exists like in the GS case.

Now, the grand canonical potential is

$$\begin{aligned} \Omega_\beta^0[v] &= \min_n \left\{ T_\beta[n] + \langle v, n \rangle \right\} = \\ &= \min_{\substack{\{p_k, \varphi_k\} \\ 0 \leq p_k \leq 1 \\ (\varphi_k | \varphi_{k'}) = \delta_{kk'}}} \left\{ \sum_k \mathcal{T}_\beta[p_k, \varphi_k] + \langle v, n[p_k, \varphi_k] \rangle \right\} \end{aligned} \quad (39)$$

where again the φ_k must be orthonormal.

Variation of the φ_k^* under the last constraint yields for the minimizing orbitals φ_k^0

$$(\hat{t} + v) \varphi_k^0 = \varphi_k^0 \varepsilon_k^0, \quad (40)$$

and variation of the p_k yields

$$p_k^0(\beta) = f_\beta(\varepsilon_k^0) = \frac{1}{e^{\beta \varepsilon_k^0} + 1}, \quad (41)$$

which is the correct result in this physically trivial case. For any $v \in X^*$ the minimum of (39) does indeed exist, and the minimizing density is

$$n_\beta^0[v] = n_{\beta ss'}^0(\mathbf{r}) = \sum_k f_\beta(\varepsilon_k^0) \varphi_k^0(\mathbf{r}, s) \varphi_k^{0*}(\mathbf{r}, s') \quad (42)$$

so that $N = \sum_k f_\beta(\varepsilon_k^0)$ relates the average particle number N to the chemical potential μ . Only the occupation numbers depend on temperature $(k\beta)^{-1}$ and on the value of the chemical potential μ from which v and the ε_k^0 are measured.

Here, $n_\beta^0 > 0$ everywhere is intuitively clear because $f_\beta(\varepsilon_k^0) > 0$ for all k .

For densities minimizing (39) it obviously holds that

$$T_\beta[n] = \Phi_\beta^0[n] = F_\beta^0[n], \quad n \in D_\beta^0. \quad (43)$$

($T_\beta[n]$ replaces the density matrix functional $T_{\text{DM}}[n]$ of the ground state theory, Eq. (9).)

Accounting for the Coulomb interaction of the electrons in mean-field approximation simply means to replace v in the above Schrödinger equation by $v + v^{\text{H}}$ where

$$v^{\text{H}}(\mathbf{r}) = \int \frac{n_{\beta s' s'}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{x}' \quad (44)$$

contains self-interaction. Since⁴ $\nabla \varphi_k \in \mathbf{L}^2(\mathbb{T}^3)$ implies $|\varphi_k|^2 \in \mathbf{L}^3(\mathbb{T}^3)$, taken as a KS ansatz $n = \sum_k p_k |\varphi_k|^2 \geq 0$ is sufficiently general for the density of an interacting system too. Densities of this type apparently form a convex domain D of the functional space $X = \mathbf{L}^3(\mathbb{T}^3)$ on which $T_\beta[n]$ is also defined by (38). ($\mathcal{J}_N \subset D$ for every real N , $0 \leq N < \infty$.)

As $D_\beta \subset D$ also for $\hat{W} \neq 0$, by

$$F_\beta[n] = T_\beta[n] + \frac{1}{2} \int \frac{n_{ss}(\mathbf{r}) n_{s' s'}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{x} d\mathbf{x}' + F_\beta^{\text{XC}}[n] \quad (45)$$

for $n \in D$ an exchange and correlation density functional $F_\beta^{\text{XC}}[n]$ is defined (since the other density functionals of this relation were previously defined or are explicitly given on D). Inserting here $n = \sum_k p_k |\varphi_k|^2$ transforms (31) into a minimum search by varying φ_k^* and p_k as above in the GS theory. The derivatives with respect to φ_k^* and p_k of F_β exist on the basis of (35) for $\{p_k, \varphi_k\}$ yielding $n \in D_\beta$, and those of the second term on the right hand side of (45) are explicitly known. Hence, the situation with T_β and F_β^{XC} is like in the GS theory. We cannot expect $D_\beta \subset D_\beta^0$.

Formally, like in (14) one may again assume

$$\begin{aligned} T_\beta[n] + F_\beta^{\text{XC}}[n] &= \\ \min_{\substack{\{p_k, \varphi_k\} \\ 0 \leq p_k \leq 1 \\ (\varphi_k | \varphi_{k'}) = \delta_{kk'}}} \left\{ \sum_k \mathcal{T}_\beta[p_k, \varphi_k] + \mathcal{F}_\beta^{\text{XC}}[\{p_k, \varphi_k\}] \mid \right. & \\ \left. \sum_k p_k |\varphi_k|^2 = n \right\} & \end{aligned} \quad (46)$$

which yields the KS equation

$$(\hat{t} + \hat{v}^{\text{eff}} - \varepsilon_k) \varphi_k = 0, \quad (47)$$

with

$$\hat{v}^{\text{eff}} \varphi_k = (v + v^{\text{H}} + \hat{v}^{\text{XC}}) \varphi_k, \quad \hat{v}^{\text{XC}} \varphi_k = \frac{1}{p_k} \frac{\delta \mathcal{F}_\beta^{\text{XC}}}{\delta \varphi_k^*}, \quad (48)$$

and $p_k(\beta) = f_\beta$ from (41) with ε_k^0 replaced by ε_k . Compare also the previous discussion of the property (15) of $\hat{v}^{\text{XC}}$. Note that we did again not prove the existence of $\delta F_\beta^{\text{XC}}/\delta n$: v^{eff} , if it exists at all, need not exist as an orbital independent local potential, it might be non-local and orbital dependent. In this respect the situation is the same as for the ground state theory.

Given the external potential v , the solutions of this KS equation determine, via the analogues of (41, 42) without superscripts, the density $n_\beta[v] \in D_\beta$ minimizing the right hand side of (31) and hence providing the grand canonical potential

$$\Omega_\beta[v - \mu] = F_\beta[n_\beta[v - \mu]] + \int (v - \mu)n_\beta[v - \mu] dx. \quad (49)$$

where we explicitly reinserted the chemical potential μ . The latter is related to the particle number N by

$$-\frac{\partial \Omega}{\partial \mu} = N = \sum_k f_\beta(\varepsilon_k - \mu) \quad (50)$$

which is also confirmed by inserting (35) and the KS expression for n into (49).

The whole theory, of course, as in the ground state variant again depends on the knowledge of the density functional $F_\beta^{\text{XC}}[n]$ and of $\mathcal{F}_\beta^{\text{XC}}$ in the KS theory, both of which are hardly ever accessible (if the latter exists at all) and hence have to be modeled changing the exact theory into a model theory within a (nearly) rigorous frame. An early *ad-hoc* application is¹³. Since (50) rests on (35), it can be used as a check for the quality of a model $F_\beta^{\text{XC}}[n]$, for instance down to which temperature it can reasonably be used for a specific answer.

Acknowledgments

I am grateful to K. Koepernik for helpful discussions.

* Electronic address: h.eschrig@ifw-dresden.de;
URL: <http://www.ifw-dresden.de/~helmut>

¹ P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964).

² W. Kohn and L. Sham, Phys. Rev. **140** A1133 (1965).

³ M. Levy, Phys. Rev. A **26**, 1200 (1982).

⁴ E. H. Lieb, Int. J. Quant. Chem. **XXIV**, 243 (1983).

⁵ N. D. Mermin, Phys. Rev. **137**, A1441 (1965).

⁶ E. g. M. V. Stoitsov and I. Zh. Petkov, Ann. Phys. **184**, 121 (1988).

⁷ R. M. Wentzcovitch, J. L. Martins and P. B. Allen, Phys. Rev. B **45**, 11372 (1992); M. Weinert and J. W. Davenport, Phys. Rev. B **45**, 13709 (1992).

⁸ For a typical example see: W. Kohn and J. M. Luttinger,

Phys. Rev. **118**, 41 (1960).

⁹ H. Eschrig and W. E. Pickett, Solid St. Commun. **118**, 123 (2001).

¹⁰ W. Kohn, A. Savin and C. A. Ullrich, Int. J. Quant. Chem. **100**, 20 (2004).

¹¹ H. Eschrig, *The Fundamentals of Density Functional theory*, Edition am Gutenbergplatz Leipzig, 2003, www.eagle-leipzig.de.

¹² see e.g. L. D. Landau and E. M. Lifshits, *Statistical Physics, Part I*, §53, Pergamon Press, London 1980.

¹³ A. Ghazali and P. Leroux Hugon, Phys. Rev. Lett. **41**, 1569 (1978).